



- ❖ Nature Genetics, 2013. Genome-wide association study identifies loci influencing concentrations of liver enzymes in plasma. J.C. Chambers et al.
- ❖ Hepatology, 2010. A Common Variant in the Patatin-Like Phospholipase 3 Gene (PNPLA3) Is Associated with Fatty Liver Disease in Obese Children and Adolescents. N. Santoro et al.
- ❖ Hepatology and Nutrition, 2014. A 4-Polymorphism Risk Score Predicts Steatohepatitis in Children With Nonalcoholic Fatty Liver Disease. V. Nobili et al.
- ❖ World Journal Gastroenterology, 2018. Promoting genetics in non-alcoholic fatty liver disease: Combined risk score through polymorphisms and clinical variables. U. Vespasiani et al.
- ❖ Hepatology, 2012. Glucokinase links Krüppel-like factor 6 to the regulation of hepatic insulin sensitivity in non-alcoholic fatty liver disease. L.P. Bechmann et al.
- ❖ Hepatology, 2012. Lower serum hepcidin and greater parenchymal iron in nonalcoholic fatty liver disease patients with C282Y HFE mutations. J. E. Nelson et al.
- ❖ Canadian Journal of Gastroenterology and Hepatology, 2018. Combining Genetic Variants to Improve Risk Prediction for NAFLD and Its Progression to Cirrhosis: A Proof of Concept Study. U. Vespasiani et al.
- ❖ European Review for Medical and Pharmacological Sciences, 2016. Antioxidant diet and genotyping as tools for the prevention of liver disease. R. Di Francia et al.